

# Complete Set of Commuting Symmetry Operators for the Klein-Gordon Equation in Generalized Higher-Dimensional Kerr-NUT-(A)dS Spacetimes

Artur Sergyeyev\*

*Mathematical Institute, Silesian University in Opava,  
Na Rybníčku 1, 74601 Opava, Czech Republic*

Pavel Krtouš†

*Institute of Theoretical Physics,  
Faculty of Mathematics and Physics,  
Charles University in Prague,  
V Holešovičkách 2, Prague, Czech Republic*

(Dated: January 7, 2008)

We consider the Klein-Gordon equation in generalized higher-dimensional Kerr-NUT-(A)dS spacetime without imposing any restrictions on the functional parameters characterizing the metric. We establish commutativity of the second-order operators constructed from the Killing tensors found in [J. High Energy Phys. **02** (2007) 004] and show that these operators, along with the first-order operators originating from the Killing vectors, form a complete set of commuting symmetry operators (i.e., integrals of motion) for the Klein-Gordon equation. Moreover, we demonstrate that the separated solutions of the Klein-Gordon equation obtained in [J. High Energy Phys. **02** (2007) 005] are joint eigenfunctions for all of these operators. We also present explicit form of the zero mode for the Klein-Gordon equation with zero mass.

In the semiclassical approximation we find that the separated solutions of the Hamilton-Jacobi equation for geodesic motion are also solutions for a set of Hamilton-Jacobi-type equations which correspond to the quadratic conserved quantities arising from the above Killing tensors.

PACS numbers: 04.50.-h, 04.50.Gh, 04.70.Bw, 04.20.Jb

## I. INTRODUCTION

Investigation of the properties of higher-dimensional black-hole spacetimes has recently attracted considerable attention, in particular in connection with the string theory. The metrics describing black holes of increasing generality were found in [1, 2, 3, 4, 5]. The most general metric of this kind known so far corresponds to a higher-dimensional generally rotating (however neither charged nor accelerated) black hole with the NUT parameters and arbitrary cosmological constant. This metric was found by Chen, Lü and Pope [6] in the form which generalizes Carter's four-dimensional Kerr-NUT-(anti-)de Sitter metric [7, 8].

The spacetime with the metric from [6] has a lot of interesting properties. In  $D$  dimensions it possesses explicit and hidden symmetries encoded in the series of  $n = [D/2]$  rank-two Killing tensors and  $D - n$  Killing vectors. The former ones can be constructed from the so-called principal Killing-Yano tensor [9], and in fact the spacetime in question is the only one admitting a rank-two closed conformal Killing-Yano tensor with certain further properties [10]. The symmetries allow one to define a complete set of  $D$  quantities conserved along geodesics. These quantities are linear and quadratic in canonical momenta. Moreover, they are functionally independent and in involution [11, 12] and thus their existence guarantees complete integrability of the geodesic motion.

lution [11, 12] and thus their existence guarantees complete integrability of the geodesic motion.

The existence of such integrals of motion is intimately related to separability of the Hamilton-Jacobi and Klein-Gordon equations. In [13] it was shown that the presence of these integrals yields the so-called separability structure. The latter guarantees separability of the Hamilton-Jacobi equation and, for the Einstein spaces, also separability of the Klein-Gordon equation. Separability of the latter equation in the spacetime under study was explicitly demonstrated in [14].

In the present paper we discuss operator counterparts of the conserved quantities constructed from the Killing vectors and rank-two Killing tensors. Namely, we convert the integrals of motion into operators using the rule  $\mathbf{p} \rightarrow -i\alpha\nabla$  and employing the symmetric ordering of derivatives, and we demonstrate that all these operators commute. Since one of these operators is, up to an overall constant factor, the Klein-Gordon operator itself, we thus obtain symmetry operators for the Klein-Gordon equation in the sense of [15, 16]. Moreover, we show that the separated solutions of the Klein-Gordon equation found in [14] are joint eigenfunctions of all symmetry operators with eigenvalues corresponding to the separation constants. As a byproduct, we obtain a zero mode solution (30) for the Klein-Gordon equation with zero mass.

We further demonstrate that semiclassical approximations of the eigenvalue equations yield a set of Hamilton-Jacobi-type equations. The latter can be solved using the separation of variables in the same fashion as in [14].

\*Electronic address: Artur.Sergyeyev@math.slu.cz

†Electronic address: Pavel.Krtous@utf.mff.cuni.cz

It is worth noticing that all these properties actually hold for a broader class of spacetimes than just the black-hole spacetimes of [6]. These properties depend on the algebraic structure of the metric (1) rather than on the explicit form of metric functions  $X_\mu$ . For this reason in what follows we do not require our metric to satisfy the vacuum Einstein equation.

## II. PRELIMINARIES

Consider  $D$ -dimensional spacetime with the metric

$$\begin{aligned} g &= \sum_{\mu=1}^n \left[ \frac{U_\mu}{X_\mu} dx_\mu^2 + \frac{X_\mu}{U_\mu} \left( \sum_{j=0}^{n-1} A_\mu^{(j)} d\psi_j \right)^2 \right] \\ &\quad + \varepsilon \frac{c}{A^{(n)}} \left( \sum_{k=0}^n A^{(k)} d\psi_k \right)^2 \\ &= \sum_{\mu=1}^n \left( \frac{U_\mu}{X_\mu} \epsilon^\mu \epsilon^\mu + \frac{X_\mu}{U_\mu} \epsilon^{\hat{\mu}} \epsilon^{\hat{\mu}} \right) + \varepsilon \frac{c}{A^{(n)}} \epsilon^{\hat{0}} \epsilon^{\hat{0}}. \end{aligned} \quad (1)$$

Here  $n = [D/2]$ ,  $\varepsilon = D - 2n$ , and  $c$  is an arbitrary constant;  $x_\mu$ ,  $\mu = 1, \dots, n$ , correspond to radial and latitudinal directions while  $\psi_k$ ,  $k = 0, \dots, n + \varepsilon - 1$ , correspond to temporal and longitudinal directions. The radial coordinate and some other related quantities are actually rescaled by the imaginary unit  $i$  in order to bring the metric into a more symmetric and compact form, cf. e.g. [6]. The signature of the metric depends on the signs of the metric functions; for the physically relevant ranges of coordinates it is  $(- + \dots +)$ .

We use Latin indices from the beginning of the alphabet to label the whole coordinate set:  $\{x^a\} = \{x_\mu, \psi_k\}$ , where  $a = 1, \dots, D$ ,  $\mu = 1, \dots, n$ , and  $k = 0, \dots, n + \varepsilon - 1$ . The non-normalized one-forms  $\{\epsilon_\mu, \epsilon_{\hat{\mu}}\}$  that diagonalize the metric, and the dual vector frame  $\{\epsilon^\mu, \epsilon^{\hat{\mu}}\}$  have the form

$$\begin{aligned} \epsilon^\mu &= dx_\mu, \quad \epsilon^{\hat{\mu}} = \sum_{j=0}^{n-1} A_\mu^{(j)} d\psi_j, \quad \epsilon^{\hat{0}} = \sum_{k=0}^n A^{(k)} d\psi_k, \\ \epsilon_\mu &= \partial_{x_\mu}, \quad \epsilon_{\hat{\mu}} = \sum_{k=0}^{n+\varepsilon-1} \frac{(-x_\mu^2)^{n-1-k}}{U_\mu} \partial_{\psi_k}, \quad \epsilon_{\hat{0}} = \frac{1}{A^{(n)}} \partial_{\psi_n}. \end{aligned} \quad (2)$$

The quantities  $\epsilon^{\hat{0}}$  and  $\epsilon_{\hat{0}}$  are defined only for odd  $D = 2n+1$ . By  $\partial_{x_\mu}$  and  $\partial_{\psi_k}$  we denote the coordinate vectors.

The functions  $U_\mu$ ,  $A_\mu^{(k)}$ ,  $U$ , and  $A^{(k)}$  used below are defined as follows

$$\begin{aligned} A_\mu^{(k)} &= \sum_{\substack{\nu_1, \dots, \nu_k=1 \\ \nu_1 < \dots < \nu_k, \nu_i \neq \mu}}^n x_{\nu_1}^2 \cdots x_{\nu_k}^2, \quad A^{(k)} = \sum_{\substack{\nu_1, \dots, \nu_k=1 \\ \nu_1 < \dots < \nu_k}}^n x_{\nu_1}^2 \cdots x_{\nu_k}^2, \\ U_\mu &= \prod_{\substack{\nu=1 \\ \nu \neq \mu}}^n (x_\nu^2 - x_\mu^2), \quad U = \prod_{\substack{\mu, \nu=1 \\ \mu < \nu}}^n (x_\mu^2 - x_\nu^2) = \det A_\mu^{(j)}. \end{aligned} \quad (3)$$

These functions satisfy the following important relations (see e.g. [14])

$$\begin{aligned} \sum_{\mu=1}^n A_\mu^{(i)} \frac{(-x_\mu^2)^{n-1-j}}{U_\mu} &= \delta_j^i, \quad \sum_{j=0}^{n-1} A_\mu^{(j)} \frac{(-x_\mu^2)^{n-1-j}}{U_\mu} = \delta_\mu^\nu, \\ \sum_{\mu=1}^n \frac{A_\mu^{(j)}}{x_\mu^2 U_\mu} &= \frac{A^{(j)}}{A^{(n)}}, \quad \sum_{j=0}^n A^{(j)} (-x_\mu^2)^{n-1-j} = 0, \end{aligned} \quad (4)$$

for  $i, j = 0, \dots, n-1$  and  $\mu, \nu = 1, \dots, n$ . The determinant of the metric in the coordinates  $\{x_\mu, \psi_k\}$  reads

$$\mathfrak{g} = \det g_{ab} = (cA^{(n)})^\varepsilon U^2. \quad (5)$$

In the above definitions we did not specify explicit form of the functions  $X_\mu$ . In what follows we just assume that  $X_\mu = X_\mu(x_\mu)$ ,  $\mu = 1, \dots, n$ , i.e., each  $X_\mu$  is a function of a single variable  $x_\mu$ . In general, the metric (1) then does not satisfy the vacuum Einstein equations. We would recover the vacuum black-hole spacetime [6, 17] by setting

$$X_\mu = b_\mu x_\mu^{1-\varepsilon} - \varepsilon \frac{c}{x_\mu^2} + \sum_{k=\varepsilon}^n c_k x_\mu^{2k}. \quad (6)$$

The constants  $c$ ,  $c_k$  and  $b_\mu$  are then related to the cosmological constant, angular momenta, mass, and NUT charges.

The spacetime with the metric (1) possesses explicit symmetries given by the Killing vectors

$$\mathbf{L}_{(k)} = \partial_{\psi_k}, \quad k = 0, \dots, n + \varepsilon - 1, \quad (7)$$

and hidden symmetries that can be generated using the principal Killing-Yano tensor [9, 18]. Namely, it was shown in [9] that (1) admits a series of rank-two Killing tensors  $\mathbf{K}_{(j)}$ ,  $j = 0, \dots, n-1$  which are diagonal in the frame (2) and read

$$\mathbf{K}_{(j)} = \sum_{\mu=1}^n A_\mu^{(j)} \left( \frac{X_\mu}{U_\mu} \epsilon_\mu \epsilon_\mu + \frac{U_\mu}{X_\mu} \epsilon_{\hat{\mu}} \epsilon_{\hat{\mu}} \right) + \varepsilon A^{(j)} \frac{A^{(n)}}{c} \epsilon_{\hat{0}} \epsilon_{\hat{0}}. \quad (8)$$

In particular, for  $j = 0$  we have  $\mathbf{K}_{(0)} = \mathbf{g}^{-1}$ , where  $\mathbf{g}^{-1}$  is the contravariant metric. We use the Killing tensors with contravariant indices as they are more convenient for the construction of conserved quantities and symmetry operators used below.

Let  $\mathcal{M}$  be our spacetime with the metric (1) and  $\mathcal{T}^*\mathcal{M}$  be the corresponding cotangent bundle. The latter is naturally endowed with the canonical Poisson bracket. Let  $\mathbf{p}$  be the one-form of canonical momenta, so that the components  $p_a$  of  $\mathbf{p}$  are canonically conjugate to  $x^a$ ,  $a = 1, \dots, D$ . Then the above Killing vectors and tensors generate conserved quantities for the geodesic motion on  $\mathcal{T}^*\mathcal{M}$ . The conserved quantities in question are linear and quadratic in  $\mathbf{p}$  and read

$$\begin{aligned} l_{(k)} &= \mathbf{L}_{(k)} \cdot \mathbf{p}, \quad k = 0, \dots, n + \varepsilon - 1, \\ k_{(j)} &= \mathbf{p} \cdot \mathbf{K}_{(j)} \cdot \mathbf{p}, \quad j = 0, \dots, n-1. \end{aligned} \quad (9)$$

Here and below  $\cdot$  denotes contraction.

It was proved in [12] that these quantities are functionally independent and in involution with respect to the canonical Poisson bracket, and hence the geodesic motion on  $\mathcal{T}^*\mathcal{M}$  is completely integrable.

Recall that the geodesic motion in this context is generated by the Hamiltonian  $H = \frac{1}{2} k_{(0)} = \frac{1}{2} \mathbf{p} \cdot \mathbf{g}^{-1} \cdot \mathbf{p}$ , and the fact that  $l_{(k)}$  and  $k_{(j)}$  are conserved quantities (i.e., integrals) for the geodesic motion means that  $l_{(k)}$  and  $k_{(j)}$  Poisson commute with  $H$ .

### III. SYMMETRY OPERATORS AND SEPARABILITY FOR THE KLEIN-GORDON EQUATION

It is natural to study the operators obtained from the above conserved quantities using the heuristic rule  $\mathbf{p} \rightarrow -i\alpha \nabla$ , where  $\nabla$  is the usual covariant derivative with respect to the metric  $\mathbf{g}$ . Upon fixing the symmetric operator ordering for the second order operators we define

$$\begin{aligned} \mathcal{L}_{(k)} &= -i\alpha \mathbf{L}_{(k)} \cdot \nabla, \\ \mathcal{K}_{(j)} &= -\alpha^2 \nabla \cdot [\mathbf{K}_{(j)} \cdot \nabla]. \end{aligned} \quad (10)$$

We employ here the convention that the square brackets do not prevent the action of the derivatives to the right. Here  $\alpha$  is a constant giving a scale to be used in order to obtain a semiclassical (geometric-optical) approximation in the next section. Of course, in the quantum context  $\alpha$  would be nothing but the Planck constant  $\hbar$ . Writing out the above operators in the coordinates  $\{x_\mu, \psi_k\}$  we obtain

$$\mathcal{L}_{(k)} = -i\alpha \frac{\partial}{\partial \psi_k}, \quad k = 0, \dots, n + \varepsilon - 1, \quad (11)$$

$$\begin{aligned} \mathcal{K}_{(j)} &= -\alpha^2 \sum_{\mu=1}^n \frac{A_\mu^{(j)}}{U_\mu} \left[ \frac{\partial}{\partial x_\mu} \left[ X_\mu \frac{\partial}{\partial x_\mu} \right] + \varepsilon \frac{X_\mu}{x_\mu} \frac{\partial}{\partial x_\mu} \right. \\ &\quad \left. + \frac{1}{X_\mu} \left[ \sum_{k=0}^{n+\varepsilon-1} (-x_\mu^2)^{n-1-k} \frac{\partial}{\partial \psi_k} \right]^2 \right] \\ &\quad - \varepsilon \frac{\alpha^2 A^{(j)}}{cA^{(n)}} \left[ \frac{\partial}{\partial \psi_n} \right]^2, \quad j = 0, \dots, n-1. \end{aligned} \quad (12)$$

Here we used the fact that  $\mathcal{K}_{(j)}$  from (10) can be written as  $\mathcal{K}_{(j)} = -\alpha^2 \mathbf{g}^{-1/2} \partial_a [\mathbf{g}^{1/2} K_{(j)}^{ab} \partial_b]$ , eq. (5), and a trivial identity  $\frac{\partial}{\partial x_\mu} (A_\mu^{(j)} U/U_\mu) = 0$ .

In [9] it was shown that the ‘quasiclassical limits’  $l_{(k)}$  and  $k_{(j)}$  of  $\mathcal{L}_{(k)}$  and  $\mathcal{K}_{(j)}$  are in involution, i.e., they Poisson commute. However, the argument of [9] does not directly imply the commutativity of the corresponding ‘quantum’ operators, i.e., of  $\mathcal{L}_{(k)}$  and  $\mathcal{K}_{(j)}$ .

Our goal is to show that the operators  $\mathcal{L}_{(k)}$  and  $\mathcal{K}_{(j)}$  commute for all  $k = 0, \dots, n + \varepsilon - 1$  and  $j = 0, \dots, n - 1$ .

The commutators of  $\mathcal{L}_{(k)}$  among themselves and of  $\mathcal{L}_{(k)}$  with  $\mathcal{K}_{(j)}$  obviously vanish, i.e.,

$$[\mathcal{L}_{(k)}, \mathcal{L}_{(l)}] = 0, \quad [\mathcal{L}_{(k)}, \mathcal{K}_{(j)}] = 0 \quad (13)$$

for all  $k, l = 0, \dots, n + \varepsilon - 1$  and  $j = 0, \dots, n - 1$ , and we only have to prove that

$$[\mathcal{K}_{(i)}, \mathcal{K}_{(j)}] = 0 \quad (14)$$

for all  $i, j = 0, \dots, n - 1$ .

To this end we first observe that the operators  $\mathcal{K}_{(j)}$  can be written as

$$\mathcal{K}_{(j)} = \sum_{\mu=1}^n \frac{A_\mu^{(j)}}{U_\mu} \tilde{\mathcal{K}}_{(\mu)}, \quad (15)$$

where

$$\begin{aligned} \tilde{\mathcal{K}}_{(\mu)} &= -\alpha^2 \left[ \frac{\partial}{\partial x_\mu} \left[ X_\mu \frac{\partial}{\partial x_\mu} \right] + \frac{\varepsilon X_\mu}{x_\mu} \frac{\partial}{\partial x_\mu} + \frac{\varepsilon}{c x_\mu^2} \left[ \frac{\partial}{\partial \psi_n} \right]^2 \right. \\ &\quad \left. + \frac{1}{X_\mu} \left[ \sum_{k=0}^{n+\varepsilon-1} (-x_\mu^2)^{n-1-k} \frac{\partial}{\partial \psi_k} \right]^2 \right]. \end{aligned} \quad (16)$$

The operators (16) enjoy a remarkable property: for any given  $\mu$  the operator  $\tilde{\mathcal{K}}_{(\mu)}$  involves only  $\partial/\partial x_\mu$  and  $x_\mu$  but does not involve  $\partial/\partial x_\nu$  and  $x_\nu$  for  $\nu \neq \mu$ . Therefore  $\tilde{\mathcal{K}}_{(\mu)}$  commute, i.e., we have

$$[\tilde{\mathcal{K}}_{(\mu)}, \tilde{\mathcal{K}}_{(\nu)}] = 0 \quad (17)$$

for all  $\mu, \nu = 1, \dots, n$ .

Using the identities (4) we find that

$$\tilde{\mathcal{K}}_{(\mu)} = \sum_{k=0}^{n-1} (-x_\mu^2)^{n-1-k} \mathcal{K}_{(k)}, \quad (18)$$

and hence the commutativity of  $\tilde{\mathcal{K}}_{(\mu)}$  entails that of  $\mathcal{K}_{(j)}$ .

Indeed, consider (17) for  $\mu \neq \nu$  and rewrite it as

$$\tilde{\mathcal{K}}_{(\mu)} \tilde{\mathcal{K}}_{(\nu)} = \tilde{\mathcal{K}}_{(\nu)} \tilde{\mathcal{K}}_{(\mu)}. \quad (19)$$

Using (18) for  $\tilde{\mathcal{K}}_{(\nu)}$  on the left-hand side of (19) and for  $\tilde{\mathcal{K}}_{(\mu)}$  on the right-hand side of (19) and employing a trivial identity  $[\tilde{\mathcal{K}}_{(\mu)}, (-x_\nu^2)^{n-1-l}] = 0$  valid for  $\mu \neq \nu$  yields

$$\sum_{l=0}^{n-1} (-x_\nu^2)^{n-1-l} \tilde{\mathcal{K}}_{(\mu)} \mathcal{K}_{(l)} = \sum_{k=0}^{n-1} (-x_\mu^2)^{n-1-k} \tilde{\mathcal{K}}_{(\nu)} \mathcal{K}_{(k)}. \quad (20)$$

Again using (18) we see that (20) boils down to

$$\sum_{k,l=0}^{n-1} (-x_\mu^2)^{n-1-k} (-x_\nu^2)^{n-1-l} [\mathcal{K}_{(k)}, \mathcal{K}_{(l)}] = 0, \quad (21)$$

whence by non-singularity of the matrix  $B_\mu^j = (-x_\mu^2)^{n-1-j}$  we readily obtain (14). It is important to stress that the

above reasoning makes substantial use of the fact that the matrix  $B_\mu^j$  is the Stäckel matrix, i.e., its  $\mu$ -th column depends on  $x_\mu$  alone, cf. e.g. [19].

Notice an important corollary of (14): since the Klein-Gordon equation  $\square\phi - \frac{m^2}{\alpha^2}\phi = 0$  can be written as  $\mathcal{K}_{(0)}\phi = -m^2\phi$ , equations (13) and (14) with  $i = 0$  imply that  $\mathcal{L}_{(k)}$  and  $\mathcal{K}_{(j)}$  are *symmetry operators* for the Klein-Gordon equation, see e.g. [15, 16] and references therein for the general theory of such operators. Let us mention that for the special case of the Kerr and Kerr–Newman metrics with  $D = 4$  this was established by Carter [20].

Relations (13) and (14) suggest that we may seek for the joint eigenfunctions  $\phi$  of the operators  $\mathcal{L}_{(k)}$  and  $\mathcal{K}_{(j)}$  with the respective eigenvalues  $\Psi_k$  and  $\Xi_j$ ,

$$\mathcal{L}_{(k)}\phi = \Psi_k\phi, \quad (22)$$

$$\mathcal{K}_{(j)}\phi = \Xi_j\phi. \quad (23)$$

We will now show that these eigenfunctions can be found by separation of variables, i.e., by assuming that they have the form (see equation (4.2) of [14])

$$\phi = \prod_{\mu=1}^n R_\mu(x_\mu) \prod_{k=0}^{n+\varepsilon-1} \exp\left(\frac{i}{\alpha}\Psi_k\psi_k\right), \quad (24)$$

with each function  $R_\mu(x_\mu)$  depending on a single variable  $x_\mu$  only.

The functions (24) clearly satisfy the equations (22). Using (18) we can combine equations (23) into an equivalent set of equations

$$\left[\tilde{\mathcal{K}}_{(\mu)} - \sum_{j=0}^{n-1} \Xi_j(-x_\mu^2)^{n-1-j}\right]\phi = 0. \quad (25)$$

Note that these equations have the same form for different  $\mu$ , i.e., the  $\mu$ -th equation (25) is obtained from the  $\nu$ -th one by replacing  $x_\nu$  by  $x_\mu$ , and vice versa. Substituting (16) into (25) we find that (24) solves all of the equations (25)—and therefore equations (23) as well—provided the functions  $R_\mu$  satisfy the following *ordinary* differential equations:

$$(X_\mu R'_\mu)' + \varepsilon \frac{X_\mu}{x_\mu} R'_\mu + \frac{1}{\alpha^2} \left( \tilde{\Xi}_\mu - \frac{\tilde{\Psi}_\mu^2}{X_\mu} \right) R_\mu = 0, \quad (26)$$

where

$$\tilde{\Psi}_\mu = \sum_{k=0}^{n+\varepsilon-1} \Psi_k (-x_\mu^2)^{n-1-k}, \quad \tilde{\Xi}_\mu = \sum_{k=0}^{n+\varepsilon-1} \Xi_k (-x_\mu^2)^{n-1-k}, \quad (27)$$

and for odd  $D = 2n+1$  we set  $\Xi_n = \Psi_n^2/c$  for convenience. Upon setting  $\alpha = 1$  in these equations we recover, of course, the separated ODEs (4.10) from [14].

Thus, we proved that the functions  $\phi$  of the form (24) with  $R_\mu$  satisfying (26) are eigenfunctions of the operators  $\mathcal{L}_{(k)}$  and  $\mathcal{K}_{(j)}$ . In particular, as  $\mathcal{K}_{(0)} = -\alpha^2\square$  (see

above), we recover the result of [14] that  $\phi$  of the form (24) satisfies the Klein-Gordon equation

$$\square\phi - \frac{m^2}{\alpha^2}\phi = 0, \quad (28)$$

where  $m^2 = -\Xi_0$ .

As a final remark note that if  $\Psi_k = 0$ ,  $k = 0, \dots, n + \varepsilon - 1$ , and  $\Xi_j = 0$ ,  $j = 0, \dots, n - 1$ , then the general solution of (26) is easily found to be

$$R_\mu(x_\mu) = h_\mu + f_\mu \int \frac{dx_\mu}{X_\mu(x_\mu)x_\mu^\varepsilon}, \quad (29)$$

where  $h_\mu$  and  $f_\mu$  are arbitrary constants. Therefore,

$$\phi_0 = \prod_{\mu=1}^n \left( h_\mu + f_\mu \int \frac{dx_\mu}{X_\mu(x_\mu)x_\mu^\varepsilon} \right) \quad (30)$$

is a zero mode, i.e., it satisfies

$$\mathcal{L}_{(j)}\phi_0 = 0, \quad \mathcal{K}_{(k)}\phi_0 = 0 \quad (31)$$

for all  $j = 0, \dots, n + \varepsilon - 1$  and  $k = 0, \dots, n - 1$ .

#### IV. HAMILTON-JACOBI EQUATION

Upon taking the solution  $\phi$  of the Klein-Gordon equation in the form

$$\phi = A \exp\left(\frac{i}{\alpha}S\right) \quad (32)$$

we find that in the semiclassical (or geometric-optical, depending on the application in question) approximation the function  $S$  satisfies the Hamilton-Jacobi equation

$$dS \cdot g^{-1} \cdot dS + m^2 = 0, \quad (33)$$

where  $m^2 = -\Xi_0$ . Recall that passing to the semiclassical approximation in our case amounts to plugging the Ansatz (32) into equation (28) multiplied by  $\alpha^2$  and taking the limit  $\alpha \rightarrow 0$ .

The same approximation leads to the Hamilton-Jacobi-type equations for each of the ‘wave’ equations (23), namely

$$dS \cdot K_{(k)} \cdot dS = \Xi_k. \quad (34)$$

In a similar fashion, the quasiclassical limit of equations (22) yields

$$L_{(k)} \cdot dS = \Psi_k. \quad (35)$$

After the above discussion of the Klein-Gordon equation it is not surprising that all these conditions are satisfied by the separated solution found in [14].

Indeed, the additive separation of variables yields [14] the following Ansatz for  $S$ :

$$S = \sum_{\mu=1}^n S_\mu(x_\mu) + \sum_{k=0}^{n+\varepsilon-1} \Psi_k\psi_k, \quad (36)$$

which automatically guarantees that the conditions (35) are satisfied.

The Hamilton-Jacobi-type equations (34) lead to the first-order differential equations

$$\sum_{\mu=1}^n \frac{A_{\mu}^{(j)}}{U_{\mu}} \left( X_{\mu} S_{\mu}'^2 + \frac{\tilde{\Psi}_{\mu}^2}{X_{\mu}} \right) + \varepsilon \frac{A^{(j)}}{cA^{(n)}} \Psi_n^2 = \Xi_j. \quad (37)$$

Taking linear combinations of these equations with the coefficients given again by the entries of the matrix  $B_{\mu}^j = (-x_{\mu}^2)^{n-1-j}$  yields an equivalent set of ordinary differential equations for  $S_{\mu}$ 's,

$$S_{\mu}'^2 = \frac{\tilde{\Xi}_{\mu}}{X_{\mu}} - \frac{\tilde{\Psi}_{\mu}^2}{X_{\mu}^2}, \quad (38)$$

where  $\tilde{\Psi}_{\mu}$  and  $\tilde{\Xi}_{\mu}$  are given by (27) and we again set  $\Xi_n = \Psi_n^2/c$  for odd  $D = 2n + 1$ . Equations (38) are precisely the separated first-order ODEs (3.7) of [14].

By direct inspection we also find that equations (38) yield the semiclassical approximation of the separability conditions (26) for the Klein-Gordon equation, with the functions  $R_{\mu}$  related to  $S_{\mu}$  as  $R_{\mu} = \exp(\frac{i}{\alpha} S_{\mu})$ .

Finally, upon identifying the momentum vector  $\mathbf{p} = dS$  in (34) and (35) we recover the original conserved quantities (9) in terms of the separation constants:

$$l_{(j)} = \Psi_j, \quad k_{(j)} = \Xi_j. \quad (39)$$

## V. CONCLUSIONS AND DISCUSSION

In the present paper we have established that the quantities  $\mathcal{L}_{(k)}$  and  $\mathcal{K}_{(j)}$ , see (10), (11) and (12), form a

complete set of commuting symmetry operators for the Klein-Gordon equation in the spacetime with the metric (1). The symmetry operators  $\mathcal{L}_{(k)}$  are associated with the Killing vectors, and the operators  $\mathcal{K}_{(j)}$  with the Killing tensors constructed from the Killing-Yano tensor [9]. We proved the commutativity of the symmetry operators in question for a general class of metrics (1) with each  $X_{\mu} = X_{\mu}(x_{\mu})$  being an arbitrary function of a single variable  $x_{\mu}$ ,  $\mu = 1, \dots, n$ ; this class includes higher-dimensional Kerr-NUT-AdS metrics [6] as special cases. We have further shown that the separated solutions (24) of the Klein-Gordon equation found in [14] provide, cf. e.g. [15, 19], joint eigenfunctions of  $\mathcal{L}_{(k)}$  and  $\mathcal{K}_{(j)}$ , see (22) and (23). We have also analyzed the quasiclassical limit of the above results and compared this limit with the results of [14]. Finally, we found explicit form of the zero mode (30) for the zero-mass Klein-Gordon equation.

In our opinion, it would be interesting to address similar issues for the Dirac equation in the spacetime with the metric (1), i.e., to find the symmetry operators whose joint eigenfunctions are the separated solutions of the Dirac equation found in [21].

## Acknowledgments

Both authors gratefully acknowledge the support from the Ministry of Education, Youth and Sports of the Czech Republic under grants MSM4781305904 (A.S.) and MSM0021610860 (P.K.). P.K. was also kindly supported by the grant GAČR 202/08/0187. The authors are pleased to thank the referee for useful suggestions. We also thank D. Kubizňák for drawing our attention to the reference [20].

- 
- [1] F. R. Tangherlini, Schwarzschild Field in  $N$  Dimensions and the Dimensionality of Space Problem, *Nuovo Cimento* **27**, 636 (1963).
  - [2] R. C. Myers and M. J. Perry, Black holes in higher dimensional space-times, *Ann. Phys. (N.Y.)* **172**, 304 (1986).
  - [3] S. W. Hawking, C. J. Hunter, and M. M. Taylor-Robinson, Rotation and the AdS/CFT correspondence, *Phys. Rev. D* **59**, 064005 (1999), arXiv:hep-th/9811056.
  - [4] G. W. Gibbons, H. Lü, D. N. Page, and C. N. Pope, Rotating Black Holes in Higher Dimensions with a Cosmological Constant, *Phys. Rev. Lett.* **93**, 171102 (2004), arXiv:hep-th/0409155.
  - [5] G. W. Gibbons, H. Lü, D. N. Page, and C. N. Pope, The General Kerr-de Sitter Metrics in All Dimensions, *J. Geom. Phys.* **53**, 49 (2005), arXiv:hep-th/0404008.
  - [6] W. Chen, H. Lü, and C. N. Pope, General Kerr-NUT-AdS metrics in all dimensions, *Class. Quantum Grav.* **23**, 5323 (2006), arXiv:hep-th/0604125.
  - [7] B. Carter, A new family of Einstein spaces, *Phys. Lett.* **26A**, 399 (1968).
  - [8] B. Carter, Hamilton-Jacobi and Schrodinger Separable Solutions of Einstein's Equations, *Commun. Math. Phys.* **10**, 280 (1968); available online at <http://projecteuclid.org/euclid.cmp/1103841118>.
  - [9] P. Krtouš, D. Kubizňák, D. N. Page, and V. P. Frolov, Killing-Yano tensors, rank-2 Killing tensors, and conserved quantities in higher dimensions, *J. High Energy Phys.* **02**, 004 (2007), arXiv:hep-th/0612029.
  - [10] T. Houri, T. Oota, and Y. Yasui, Closed conformal Killing-Yano tensor and Kerr-NUT-de Sitter spacetime uniqueness, *Phys. Lett.* **B656**, 214 (2007), arXiv:0708.1368 [hep-th].
  - [11] D. N. Page, D. Kubizňák, M. Vasudevan, and P. Krtouš, Complete Integrability of Geodesic Motion in General Higher-Dimensional Rotating Black Hole Spacetimes, *Phys. Rev. Lett.* **98**, 061102 (2007), arXiv:hep-th/0611083.
  - [12] P. Krtouš, D. Kubizňák, D. N. Page, and M. Vasudevan, Constants of Geodesic Motion in Higher-Dimensional Black-Hole Spacetimes, *Phys. Rev. D* **76**, 084034 (2007), arXiv:0707.0001 [hep-th].
  - [13] S. Benenti and M. Francaviglia, Remarks on certain separability structures and their applications to general relativity, *Gen. Rel. Grav.* **10**, 79 (1979).

- [14] V. P. Frolov, P. Krtouš, and D. Kubizňák, Separability of Hamilton-Jacobi and Klein-Gordon equations in general Kerr-NUT-AdS spacetimes, *J. High Energy Phys.* **02**, 005 (2007), arXiv:hep-th/0611245.
- [15] W. Miller, Jr, *Symmetry and Separation of Variables* (Addison-Wesley, Reading, Massachusetts, 1977); available online at <http://www.ima.umn.edu/~miller/separationofvariables.html>.
- [16] W. I. Fushchich and A. G. Nikitin, *Symmetry of Equations of Quantum Mechanics* (Allerton, New York, 1994).
- [17] N. Hamamoto, T. Houri, T. Oota, and Y. Yasui, Kerr-NUT-de Sitter curvature in all dimensions, *J. Phys.* **A40**, F177 (2007), arXiv:hep-th/0611285.
- [18] D. Kubizňák and V. P. Frolov, Hidden symmetry of higher dimensional Kerr-NUT-AdS spacetimes, *Class. Quantum Grav.* **24**, F1 (2007), arXiv:gr-qc/0610144.
- [19] E. G. Kalnins and W. Miller, Jr, The theory of orthogonal R-separation for Helmholtz equations, *Adv. Math.* **51**, 91 (1984); available online at <http://www.ima.umn.edu/~miller/Rsep.pdf>.
- [20] B. Carter, Killing tensor quantum numbers and conserved currents in curved space, *Phys. Rev. D* **16**, 3395 (1977).
- [21] T. Oota and Y. Yasui, Separability of the Dirac equation in higher dimensional Kerr-NUT-de Sitter spacetime, (2007), arXiv:0711.0078 [hep-th].